

Research Article

Studies on the Formation and Composition of Copper(II)- α - and γ -Halo-1-Oxobutanilides Complexes

M. A. Zayed

Chemistry Department, Faculty of Science, Cairo University, Giza 12613, Egypt
Address correspondence to M. A. Zayed, mazayed429@yahoo.com

Received 21 October 2023; Revised 2 December 2023; Accepted 8 December 2023

Copyright © 2023 M. A. Zayed. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract The formation, structure, and relative stabilities of copper(II) complexes with γ -bromo-, p-tolyl- γ -bromo-, γ -iodo-, α -bromo-, α -chloro-, and α -phenyl-azo- γ -bromoacetoacetanilide (L) are investigated. The ultraviolet spectrophotometric study indicates the formation of the stable complex species Cu(L)_{1-3} in solution. The formation constants of these complexes were computed from the spectrophotometric data. The solid complexes were prepared and chemically analyzed and their FT-IR spectra were studied. The structural formulae of the prepared chelates were proposed.

Keywords structure; relative stability; oxobutanilides; copper complexes, absorbing species

1. Introduction

Copper(II) complexes and other metal cations with acetoacetanilide and some of its derivatives have been studied by a group of investigators using different experimental techniques [1,2,3,4]. Zayed et al. [5] studied structures of 2-thiohydantoin and its metal chelates. Also, Zayed et al. [6] determined the stability constants of uranyl complexes potentiometrically. The effect of substituents on the phenyl ring and on the stability of these complexes was interpreted in terms of inductive, mesomeric, and steric effects. CuL_2 complexes (HL = acetoacetanilide, o-chloroacetoacetanilide, and o-acetoacetatoludide) were prepared, their spectra were studied, and their stability constants were determined [7,8]. The stability constants of 13 oximes of acetoacetanilide and Cu^{2+} , UO_2^{2+} , and Be^{2+} complexes were determined by Kettrup [9] using potentiometric pH-measurements.

The wide applications and great importance of complexes of acetoacetanilide and its derivatives with copper directed our attention toward further studies on their structures, which was the aim of the present investigation. Acetoacetanilide complex can be used as fungicides and has a great influence in nitrification of soil. The complexes of these compounds with cations, especially those of Cr^{3+} and Cu^{2+} , are used for water proofing, tanning, and coloring and as parasiticides [10].

2. Experimental

2.1. Materials and methods

All chemicals used were of the highest purity available. They included γ -bromoacetoacetanilide (γ -Br-A.A.A.), p-tolyl- γ -bromoacetoacetanilide (p-tolyl- γ -Br-A.A.A.), γ -iodoacetanilide (γ -I-A.A.A.), α -bromoacetoacetanilide (α -Br-A.A.A.), α -chloroacetanilide (α -Cl-A.A.A.), and α -phenyl-azo- γ -bromoacetoacetanilide (α - C_6H_5 -azo- γ -Br-A.A.A.) [11].

2.1.1. Preparation of complexes and their analysis

The solid complexes of copper acetate with γ - and α -haloacetoacetanilide were prepared by adding the calculated amount of ligand (dissolved in the least amount of ethanol) dropwise to 0.2–0.5 g copper acetate (dissolved in the least amount of water) with constant stirring. The solid complexes were spontaneously precipitated and separated from solution by filtration, washed with alcohol-water mixture, and left to dry in a vacuum desiccator for few days. These complexes were recrystallized from ethyl alcohol or acetone.

The molecular weights of these complexes were calculated from EDTA titration data [12]. Microanalyses of precipitated complexes for C, H, N, halogen, and metal were performed using the recommended procedures. The results are shown in Table 1.

2.1.2. Spectrophotometric studies

(I) Ultraviolet spectral measurements

The spectra of copper acetate and copper complexes with γ - and α -halo derivatives were scanned in 210–360 nm using a Beckmann DK-L recording spectrophotometer. The formation and composition of different copper complexes in solution were studied by applying Job's continuous variation method in the case of α -Br-A.A.A. and the molar ratio method (mrm) in the case of all ligands used [12].

Table 1: Analysis of copper(II) acetoactanilide complexes.

Ligand (L)	Color of the complex	Melting point		C%		H%		N%		Other elements		M. Wt.		Proposed formula of the complex M:L
		pure	mixed	Calc.	found	Calc.	found	Calc.	found	Calc.	found	Calc.	found	
γ -Br-A.A.A.	Green	C*	C*							Cu%				$(\text{CuL}_2)(\text{AcO})_2$
		164	146	41.5	42.0	3.70	3.0	4.03	4.6	9.15	8.9	693.76	694.06	
												694.85		
p-tolyl- γ -Br-A.A.A.	Faint green									Br%				$(\text{CuL}_3)(\text{AcO})_2$
		166	144		41.5		3.0		4.4					
3-phenyl-azo- γ -Br-A.A.A.	Chocolate brown													$(\text{CuL})_2(\text{AcO})_2$
α -Br-A.A.A.	Pale green													$(\text{CuL}_2)(\text{AcO})_2$
γ -I-A.A.A.	Bright green													$(\text{CuL}_2)(\text{AcO})_2$
α -Cl-A.A.A.	Green													$(\text{CuL}_2)(\text{AcO})_2$

*Molecular weights determined in the microanalytical unit.

(II) Infrared spectral measurements

The FT-IR spectra of the pure ligand and solid complexes were obtained by applying the KBr disc technique and using Pye Unicam SP1000 spectrophotometer.

3. Results and discussion

3.1. UV spectra

The spectra of copper acetate, ligands, and complex were scanned in the wavelength range of 210–360 nm. Representative curves for the data obtained by Job's method for Cu- γ -Br-A.A.A. complexes are shown in Figure 1.

The results obtained in this case are represented by the plot of absorbance against wavelength exhibiting maxima for various metals to ligand ratios, which may be attributed to the presence of more than one complex species. The presence of the most stable ML_2 complex is confirmed by the plot of absorbance against mL reagent at 300 and 330 nm. The data obtained in the case of other ligands by applying the mrm indicate the formation of the most stable 1:2 (M:L) complexes, except in the case of 3- C_6H_5 -azo- γ -Br-A.A.A. where the results revealed the formation of 1:1 and 1:2 (M:L) adducts using the previously published technique of oxygen flask combustion [13].

3.2. Determination of the number of absorbing species

In order to determine the number of absorbing species in the cases of Cu(II) complexes with the above-mentioned ligands, Coleman graphical method is applied to the data obtained by applying the mrm (Tables 2–4). By plotting the variables $x = A_{ij} - A_{i'j'}$ against $y = A_{ij} - A_{i'j'}$, where i and j are two wavelengths and i' and j' are the corresponding concentrations of the investigated solutions,




Figure 1: The ultraviolet absorption spectra of copper acetate. (a) γ -Br-A.A.A., and their complexes formed by the continuous variation method. (b) (Cu): 8×10^{-5} M, Br-A.A.A.: 9×10^{-5} M, and complexes of copper: ligand in ratios (1 = 9:1, 2 = 8:2, 3 = 7:3, 4 = 6:4, 5 = 5:5, 6 = 4:6, 7 = 3:7, 8 = 2:8, and 9 = 1:9).

straight lines were obtained, passing by the origin, in the cases of Cu- γ -Br-, p-tolyl- γ -Br-, γ -iodo-, α -Br-, and α -Cl-A.A.A., indicating the presence of two absorbing species in solution, namely, the ligand and the stable 1:2 (M:L) complex. Figure 2 is a representative of this. These are interpreted depending upon previously published work by Zayed et al. [14, 15, 16, 17, 18].

Table 2: Absorption matrices for the copper complex with acetoacetanilide derivatives in the UV range. (a) Cu- γ -Br-A.A.A., (b) Cu-p-tolyl- γ -Br-A.A.A., (c) Cu- α -phenyl-azo- γ -Br-A.A.A., and (d) Cu- γ -I-A.A.A.

		[γ -Br-A.A.A.] $\times 10^{-5}$ M							
Wavelength		1.0	2.0	3.0	4.0	5.0	6.0	7.0	8.0
		(Soln. no., j)							
i	nm	1	2	3	4	5	6	7	8
1	300	0.175	0.260	0.312	0.405	0.480	0.550	0.595	0.680
2	306	0.178	0.253	0.297	0.380	0.412	0.500	0.525	0.620
3	294	0.165	0.255	0.419	0.411	0.490	0.580	0.632	0.723
4	240	0.151	0.270	0.370	0.475	0.597	0.755	0.865	1.000
5	228	0.140	0.240	0.303	0.387	0.480	0.600	0.670	0.770

(a)

		[p-tolyl- γ -Br-A.A.A.] $\times 10^{-5}$ M							
Wavelength		1.0	2.0	3.0	4.0	5.0	6.0	7.0	8.0
		(Soln. no., j)							
i	nm	1	2	3	4	5	6	7	8
1	306	0.200	0.295	0.340	0.375	0.503	0.600	0.578	0.640
2	294	0.192	0.292	0.292	0.422	0.542	0.668	0.692	0.741
3	288	0.180	0.280	0.350	0.431	0.542	0.668	0.690	0.768
4	270	0.148	0.232	0.312	0.416	0.498	0.610	0.650	0.710
5	240	0.199	0.324	0.454	0.625	0.735	0.935	1.020	1.165

(b)

		[3-phenyl-azo- γ -Br-A.A.A.] $\times 10^{-5}$ M					
Wavelength		0.5	1.0	2.0	3.0	4.0	5.0
		(Soln. no., j)					
i	nm	1	2	3	4	5	6
1	240	0.105	0.170	0.364	0.480	0.595	7.15
2	270	0.075	0.135	0.205	0.380	0.470	0.545
3	276	0.080	0.140	0.310	0.410	0.505	0.600

(c)

		[γ -I-A.A.A.] $\times 10^{-5}$ M							
Wavelength		0.5	1.0	1.5	1.75	2.0	3.0	3.5	4.0
		(Soln. no., j)							
i	nm	1	2	3	4	5	6	7	8
1	228	0.130	0.235	0.360	0.385	0.460	0.620	0.710	0.730
2	240	0.120	0.237	0.380	0.410	0.510	0.710	0.755	0.813
3	252	0.088	0.190	0.320	0.335	0.410	0.590	0.625	0.645
4	270	0.075	0.148	0.230	0.250	0.290	0.395	0.435	0.395
5	300	0.135	0.250	0.360	0.415	0.470	0.580	0.640	0.670
6	310	0.160	0.300	0.415	0.460	0.520	0.620	0.680	0.710

(d)

Table 3: Absorption matrices for the copper complexes with acetoacetanilide derivatives in the UV range. Cu- α -Br-A.A.A.

		[α -Br-A.A.A.] $\times 10^{-5}$ M									
Wavelength		0.5	1.0	1.5	2.0	2.5	3.0	4.0	5.0	6.0	4.0
		(Soln. no., j)									
i	nm	1	2	3	4	5	6	7	8	9	10
1	288	0.052	0.058	0.090	0.105	0.130	0.150	0.185	0.230	0.270	0.290
2	252	0.108	0.152	0.203	0.252	0.295	0.355	0.445	0.540	0.640	0.690
3	246	0.104	0.148	0.195	0.235	0.275	0.335	0.410	0.505	0.625	0.695
4	240	0.108	0.138	0.158	0.215	0.245	0.293	0.360	0.422	0.501	0.535
5	228	0.115	0.140	0.195	0.218	0.265	0.300	0.345	0.392	0.400	0.415
6	288	0.052	0.058	0.090	0.105	0.130	0.150	0.185	0.230	0.270	0.290




Figure 2: Number of absorbing species in solution from spectrophotometric data in the UV range: (a) Cu- α -phenyl-azo- γ -Br-A.A.A. absorbing species; (b) Cu- γ -I-A.A.A. absorbing species.

Table 4: The calculated conventional formation constants of complexes.

Ligand	$\log K$ of 1:2 (M:L) complexes
γ -I-A.A.A.	10.50
α -Br-A.A.A.	11.13
γ -Br-A.A.A.	11.28
α -Cl-A.A.A.	11.77
3-C ₆ H ₅ -azo- γ -Br-A.A.A.	11.95
p-C ₆ H ₇ - γ -Br-A.A.A.	12.65

On the other hand, by plotting,

$$x = (A_{2j} - A_{2j'}) / (A_{1j} - A_{1j'}).$$

In the case of 3-C₆H₅-azo- γ -Br-A.A.A., a straight line is obtained revealing the existence of three absorbing species (Table 1), namely, the organic ligand and the 1:1 and 1:2 (M:L) complexes. All these complex species have been isolated in the solid state and their compositions are confirmed by chemical analysis and FT-IR spectra.

These data are interpreted depending upon the previously published data by Zayed et al. [18, 19, 20, 21] and used to prove the proposed structures of studied complexes.

3.3. Determination of the stability constants

The calculated conventional formation constants using the recommended method $\log K$ of 1:2 (M:L) are tabulated in Table 4.

These values indicate higher stabilities than those of copper-phenyl substituted acetoacetanilide (6.4–7.27 g in 50% dioxane, 8.08–9.3 g in 75% dioxane, and 5.45–10.57 g in 30–60% dioxane 1 for all possible N -substituted

anilides 2, obtained from potentiometric-pH measurements). They also indicate that stability increases in cases where there are substituents in the α -position and phenyl ring simultaneously as in the case of p-tolyl- γ -Br-A.A.A. ($\log K = 12.65$). The lower $\log K$ values obtained by potentiometric methods may be interpreted on the basis that in potentiometry, the acidity, equivalent to the protons liberated when the metal ion is chelated with the enol form of the reagent, is measured. However, since a great portion of the reagent is present in the ketoform, it is clear that the latter will contribute to the stability of the complex.

3.4. Infrared absorption spectra (FT-IR)

The FT-IR spectra of all separated solid complexes were studied. In the case of the Cu- γ -Br-A.A.A. complex, as a representative example, its spectrum (Figure 3) showed intensified broad band at 1,730 cm⁻¹ and anilide carbonyl band at 1,660 cm⁻¹. Comparing the FT-IR bands of the complex with that of the γ -Br-A.A.A., it is clear that the two carbonyls are linked via coordinate bonds to Cu(II) ions. The more intensified NH band at 3,300 cm⁻¹ in the 1:2 complexes than that in the ligand supports the presence of two moles of the ligand in the complex [14, 15, 16, 17]. The halogen carbon peaks in both ligands and metal chelates skeletons are intensively present in wave numbers 625–800 cm⁻¹; their positions are varied in ligands due to the variation of halogen kind and also in chelates as a result of complex nature and stability.

The general formula of the given complexes depending on the FT-IR studies may tentatively be represented by the equilibrium of both keto and enol forms (I and II) of the ligand attached to copper ions as given in Figure 4.




Figure 3: The infrared spectra of γ -Br-A.A.A. and its complexes with copper acetate. (a) γ -Br-A.A.A., (b) Cu-Br-A.A.A. (1:3), (c) copper acetate, and (d) Cu- γ -Br-A.A.A. (1:2).




Figure 4: The proposed equilibrium between keto-enol structures of γ -Br-A.A.A. copper complexes.

Conflict of interest The author declares that he has no conflict of interest.

References

- [1] H. Khalifa, M. M. Khater, and M. A. Zayed, *Absorption spectra of copper and palladium complexes of arylazo acetylacetone*, Egypt J Chem, 23 (1980), 311–321.
- [2] M. M. Khater, B. G. Ateya, and M. A. Zayed, *Electrochemical behavior of copper in alcoholic media. Effect of γ -haloacetoacetanilide as chelating agent*, Egypt J Chem, 26 (1983), 1–11.
- [3] M. A. Zayed, *Microdetermination of rare earth elements in their acetylpyridine N-oxide complexes by the oxine method after oxygen flask combustion*, Microchem J, 34 (1986), 200–203.
- [4] M. A. Zayed, F. A. Nour El-Dien, and M. A. F. Sharaf, *The use of thermal and spectrophotometric analyses for the structure investigation of arylhydrazonacetactylaminopyridines and their copper chelates*, Thermochem Acta, 146 (1989), 11–113.
- [5] F. A. Nour El Dien, M. A. Abdelaziz, and M. A. Zayed, *Structure identification of 2-thiohydantoin and its metal chelates by thermal and spectral analysis*, Thermochem Acta, 162 (1990), 399–407.
- [6] M. A. Zayed, A. E. Arifien, and E. A. Abou El-Nasr, *Spectral studies on uranium–pyrocatechol complexes and their applications for determination of uranium*, Egypt J Chem, 39 (1996), 157–169.
- [7] M. J. Hynes and E. M. Clarke, *Reactions of metal ions with β -ketoamides*, J Chem Soc Perkin 2, 1998 (1998), 1263–1268.
- [8] M. A. Zayed, A. E. A. Arifien, and F. S. M. Hassan, *Investigation of complexes of diphenylphosphine derivatives by thermal and other physicochemical methods of analyses*, J Therm Anal Calorim, 55 (1999), 205–212.
- [9] A. Kettrup and T. Neustadt, *3-monoximes of substituted acetoacetamides as chelating agents*, Z-Naturforsch. B, 28 (1973), 86–89.
- [10] K. N. Mehrotra, V. P. Mehta, and T. N. Nagar, *Kinetics of thermal decomposition of copper soaps*, Z-Naturforsch. B, 28 (1973), 90–92.
- [11] M. A. Zayed and H. Fischer, *Investigation of novel organometallics by thermal spectroscopic methods*, J Therm Anal Calorim, 61 (2000), 897–908.
- [12] G. G. Mohamed, M. A. Zayed, and N. E. El-Gamel, *Thermal and kinetic studies on solid complexes of 2-(2-benzimidazolylazo)-4-acetamidophenol with some transition metals*, Spectrochim Acta A Mol Biomol Spectrosc, 58 (2002), 3167–3178.
- [13] P. Borda, A. E. Crease, and P. Legzdins, *The determination of lanthanides in organometallic complexes by the closed oxygen flask method*, Microchem J, 18 (1973), 172–177.
- [14] M. A. Zayed and S. M. Abdallah, *Synthesis, characterization and electronic spectra of cefadroxil complexes of d-block elements*, Spectrochim Acta A Mol Biomol Spectrosc, 60 (2004), 2215–2224.
- [15] M. A. Zayed and S. M. Abdallah, *Synthesis and structure investigation of the antibiotic amoxicillin complexes of d-block elements*, Spectrochim Acta A Mol Biomol Spectrosc, 61 (2005), 2231–2238.

- [16] S. M. Abdallah, G. G. Mohamed, M. A. Zayed, and M. S. Abou El-Ela, *Spectroscopic study of molecular structures of novel Schiff base derived from o-phthaldehyde and 2-aminophenol and its coordination compounds together with their biological activity*, Spectrochim Acta A Mol Biomol Spectrosc, 73 (2009), 833–840.
- [17] E. M. Zayed, M. A. Zayed, and A. M. M. Hindy, *Thermal and spectroscopic investigation of novel Schiff base, its metal complexes, and their biological activities*, J Therm Anal Calorim, 116 (2014), 391–400.
- [18] E. M. Zayed, M. A. Zayed, A. M. Fahim, and F. A. El-Samahy, *Synthesis of novel macrocyclic Schiff's-base and its complexes having N_2O_2 group of donor atoms. Characterization and anticancer screening are studied*, Appl Organometal Chem, 31 (2017), e3694.
- [19] M. A. Zayed, A. Belal, and S. M. A. H. Ragha, *Structure studies of the prepared novel hydrazone Schiff's base complexes using spectroscopic, thermal analyses and their biological activities*, J Transit Met Complexes, 1 (2018), article no. 236047.
- [20] K. R. Oraby, F. S. M. Hassan, M. A. Zayed, A. E. Mohamed, and I. M. Abdellah, *DFT, TD-DFT and biological activity studies of some maleanilic acid derivatives ligands and their organometallic complexes*, Indian J Chem A, 60 (2021), 1564–1573.
- [21] M. A. Zayed, S. A. Abdel-Gawad, H. M. Abdel-Aziz, and Z. A. Abo-Ayad, *Green synthesis of nano-zero-valent copper for the D-blue 60 textile dye removal from aqueous medium*, Int J Environ Res, 17 (2023), article no. 12.